

VOC Monitoring Near the World Trade Center Site
New York State Department of Health
September 28, 2001

On September 28, 2001, New York State Department of Health (NYSDOH) and New York City Department of Health (NYCDOH) collected air samples at three locations north, southwest and southeast of the World Trade Center to measure community exposure levels to a limited number of VOCs. The locations were selected to represent the communities closest to the site that are currently occupied or might be expected to be re-occupied soon. The primary purpose was for screening, i.e. to determine whether additional VOC sampling in the surrounding community is warranted on some regular basis.

The air samples were collected in Summa canisters over two hours at three locations, and a grab sample was taken on the rubble pile (see table below). The day began sunny but became cloudy as the day progressed. A light rain started just before the evening sampling and was intermittent throughout that sampling period. Light winds were primarily from the north but shifted somewhat between the northeast and northwest throughout the day.

Location	Elevation	Time*
75 Park Place (Park Place and Greenwich St.)	roof above 14th floor 10th floor balcony street	M, A, E M, A M, A, E
Parc Place at 225 Rector	roof above 24th floor street	M, A, E M, A, E
140 Broadway (Broadway and Liberty)	roof above 52nd floor street	M, A, E M, A, E
Top of rubble		8PM**
* M - morning (~6-8AM) A - afternoon (~2-4PM) E - evening (~6-8PM)		** Grab sample (~25 seconds)

The Summa canisters were returned to NYSDOH Wadsworth Laboratories for analysis by gas chromatography – mass spectrometry (GC-MS) following EPA method TO-14.

Environmental Results and Conclusions

1. The air sample collected in the smoke plume on the WTC debris pile contained the highest levels of VOCs. VOCs exceeding 100 mcg/cubic meter were acetone and a number of fuel components or products of combustion, i.e. benzene, toluene, ethylbenzene, styrene and chloromethane. None of the reported levels were unusually high compared to samples from other fires (P.S.J. Lees, 1995). These sample results are generally consistent with similar samples of the smoke plume taken on several occasions by USEPA. However, the benzene

concentrations were not as high (in absolute terms or relative to toluene and xylene) as most of the EPA sampling results.

2. At 75 Park Place (upwind of the WTC site) and at 225 Rector (cross-wind), VOC levels, except ethanol, were in the range of levels commonly found in outdoor air samples (background). At 140 Broadway (downwind), where the smoke plume from the WTC site was visible during each of the sampling events, most of the VOCs were greater than background (generally less than five times background levels and all (except trichlorofluoromethane) less than about ten times background levels).
3. Generally, VOC concentrations at 140 Broadway were about 3 times greater than those at Park Place and Rector Place sites.
4. During each of the sampling events, the smoke plume was visibly evident at 140 Broadway and fire odors were noticeable at the street level. During sampling at the other sites, the smoke plume was not visibly evident at the sites. WTC-related odors were sometimes noticed at the Rector Place site and not noticed at the Park Place site.
5. VOC concentrations on the roofs were not substantially different from street-level VOC concentrations at each of the sampling sites, including 140 Broadway (53 stories high).
6. Concentrations of VOCs associated with the smoke plume were about 20 times greater in the grab sample at the pile than at 140 Broadway.
7. High levels of trichlorofluoromethane (a refrigerant, R-11), were observed in the samples from the roof at 140 Broadway. The building manager said that a chiller leak sensor had detected a minor refrigerant leak of R-11 on Friday (Sept 28) morning. The problem was corrected later in the day. The chiller is located on the 52nd floor and a mechanical room vent was about three meters (10 feet) from the sampler.
8. Elevated methylene chloride concentrations were also detected on the roof of 140 Broadway with slightly higher levels found in the afternoon sample when the chiller was being repaired. Methylene chloride is commonly used as a degreaser in building HVAC system maintenance and repair work. It was not measureable in the plume sample or any of the other samples.
9. The field forms for the morning samples for 140 Broadway (roof and street level) were switched so the laboratory report forms misidentify the roof and street level samples at this location.
10. Because the trichlorofluoromethane and methylene chloride were on the roof top and people are not being directly exposed to them, they are not discussed in the health implications section.
11. Assuming that the fires and smoke will continue to diminish as the recovery work progresses, these data suggest that additional VOCs sampling for these compounds is not necessary.

Data Limitations

These VOCs were selected for analysis because they are commonly found at low levels in outdoor and indoor air, routine analytical procedures were readily available and many have been shown to be constituents of fires. Other VOCs and semivolatile organic compounds have been found in the smoke from fires and can be expected near the site and in the smoke plume as well. Different sampling techniques are required for these compounds.

Public Health Implications

Although the highest levels of VOCs were found in samples from the plume at the WTC debris pile, these results are not likely to represent the potential exposure of the residents. Residents are unlikely to be in the plume at the debris pile where they could be exposed to these levels for any length of time. Therefore, the sampling results for areas potentially affected by the plume (i.e., at downwind locations) but not at the debris pile are more relevant for evaluating potential exposure to the residents. The levels of some of the chemicals in the plume measured at the debris are elevated and exceed typical background ranges, and recovery and emergency workers should take precautions to limit their exposure to emissions from the plume.

The levels of VOCs at the upwind sites (75 Park Place and 225 Rector) are within the range of typical background levels.

The air concentrations of VOCs at the downwind site (140 Broadway) were generally greater than those measured upwind (75 Park Place and 225 Rector) and typical background concentrations. However, none of the WTC-related chemicals measured at 140 Broadway exceed both typical background levels and air comparison values except benzene, chloromethane and styrene. Based on the available information, exposures to these chemicals from the plume are unlikely to last more than one year. Thus, in addition to typical background levels, the concentrations of the chemicals measured at 140 Broadway are evaluated with health comparison values for air based on noncancer effects and exposure durations of less than one year (e.g., acute (up to two weeks) and intermediate (up to one year) air comparison values).

None of the chemicals detected above background at 140 Broadway exceed intermediate or short-term air comparison values except benzene. The average and highest levels of benzene at 140 Broadway are slightly above typical background and are near the intermediate comparison value for this chemical. This comparison value for benzene is about 90 times lower than the exposure level that caused mild neurological effects in mice exposed regularly for 30 days.

The benzene air concentrations at 140 Broadway (as well as typical background concentrations of benzene) exceed the health comparison value for benzene in air based on carcinogenic effects following long-term exposure. The cancer comparison value is the air concentration associated with an increased lifetime cancer risk of one-in-one million (10^{-6}). Because the estimated duration of exposure is at most one year, and because elevated levels are unlikely to persist for extended periods of time (e.g. the fires will be put out and the wind direction will periodically

change), community exposures are not expected to result in cancer risks that are appreciably increased over those resulting from background exposure. These estimated cancer risks do not exceed one-in-ten thousand (10^{-4}), above which actions are usually taken to reduce exposure.

The average levels of chloromethane and styrene also exceed background and their air comparison values based on cancer effects and long-term exposure. Again, based on the low potential for long-term exposure to elevated levels, this is not expected to result in an appreciable increased cancer risk, but one similar to the risks resulting from exposure to background levels.

Limitations of Health Assessment

The primary limitation of this assessment is that the list of VOCs measured may not be completely representative of the emissions from the fires and recovery activities at the World Trade Center site. The available analytical methods cannot include all of the numerous combustion products that can be generated from the fires, which, in addition to the measured VOCs, could affect air quality and public health. Some chemicals that are known to be produced by fires (e.g. aldehydes, acrolein) and are strong respiratory irritants were not measured.

When odors are present, it is not unusual for some people to experience short-term effects such as respiratory irritation and effects such as headaches and dizziness. These effects tend to diminish once exposure to the odors is stopped, and usually do not result in long-term health consequences. Some people are more likely to experience these effects than others.

Conclusion

The results of the monitoring suggest that the levels of some VOCs in the plume at the debris pile are significantly elevated above typical background levels, and workers at the site should take precautions to limit their exposure. For the VOCs that were measured, the air concentrations outside the work area are not a significant health concern. However, the elevation above background levels of some VOCs indicates that combustion products may be contributing to symptoms of eye, nose and throat irritation that have been reported in workers and residents in the WTC neighborhood. Based on the current results and the expected mitigation of VOC emissions from the site in the future by controlling the fires, continued monitoring for the VOCs sampled is not necessary at this time. It may be useful to assess VOC concentrations again once the fires are controlled to ensure that other recovery activities are not a continuing source of VOCs in the community.

References

Peter S.J. Lees. 1995. Combustion products and other firefighter exposures: A state of the art review. *Occup. Med.* 10:691-706.

VOC Results 09-28.doc
Oct 5, 2001

**Summary of Outdoor Air Samples Collected at the World Trade Center Disaster Area
28 September 2001**

All Results are Parts Per Billion by Volume

Analyte	75 Park Place									Parc Place at 225 Rector									140 Broadway									N Tower Plume SPM	Outdoor Background ³		
	Roof Above 14th Floor			10th Floor			Street			Roof Above 24th Floor			Street			Roof Above 52nd Floor			Street			M	A	E	M	A	E		M	A	E
	M	A	E	M	A	E	M	A	E	M	A	E	M	A	E	M	A	E	M	A	E										
DICHLORO(DIFLUORO)METHANE	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.6	0.7	0.6	0.6	0.6	0.6	0.6	0.6	0.8	1.1	1.1	0.8	0.6	0.7	1.2				0.32 - 0.35	<0.2				
CHLOROMETHANE	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	1.0 PL	2.8	1.2	1.0 PL	3.2	2.4	1.7	58				0.6 - 0.7	0.24 - 0.63				
BROMOMETHANE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	3.9				0.05 - 3.1	<0.3				
CHLOROETHANE	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	<0.8	2.7				NA	<0.2				
ETHYL ALCOHOL	4.7	1.9	3.6	3.5	4.1	10.1	9.8	4.7	2.6	2.1	3.8	2.1	1.3	2.0	3.7	5.0	4.3	9.6	3.7	4.6	32						NA	<0.5 - 0.9			
TRICHLOROFLUOROMETHANE	0.4 PL	0.4 PL	0.4 PL	0.4 PL	0.4 PL	0.4	0.4	0.4	0.4 PL	0.4 PL	0.4 PL	0.4 PL	0.4 PL	0.4 PL	>133	>133	>133	4.8	0.6	1.6	1.7				0.19 - 0.21	<0.16					
ISOPRENE	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	2.5				NA	NA				
ACETONE	1.3	0.9	1.1	1.4	2.2	1.2	1.5	1.3	1.2	1.0	1.7	1.4	1.0	1.2	4.0	7.6	4.6	3.2	2.9	2.6	59				ND - 2.83	NA					
METHYLENE CHLORIDE	0.6 PL	<0.6	<0.6	0.6 PL	0.6 PL	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6	<0.6	2.2	0.4	3.2	<0.6	0.6 PL	<0.6	0.6 PL						0.3 - 1.8	0.14 - 1.1			
METHYL-tert-BUTYL ETHER	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	1.2	0.6 PL	0.8	0.6 PL	0.6	1.4	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6	0.6 PL	0.6 PL	0.6 PL						NA	<0.3		
n-HEXANE	0.6 PL	<0.6	<0.6	0.6 PL	0.6 PL	0.6 PL	<0.6	2.3	0.6 PL	<0.6	0.6 PL	<0.6	<0.6	<0.6	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	0.6 PL	3.7						0.83 - 2.95	0.1 - 0.5			
METHYL ETHYL KETONE	<0.7	<0.7	<0.7	<0.7	0.7 PL	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	0.7 PL	<0.7	<0.7	<0.7	1.0	1.3	0.8	0.7	0.7	0.7 PL	25					ND	NA			
TETRAHYDROFURAN	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	0.7 PL	<0.7	<0.7	<0.7	<0.7	<0.7	<0.7	3.4					NA	NA			
n-HEPTANE	<0.5	<0.5	<0.5	<0.5	0.5 PL	<0.5	0.5 PL	<0.5	<0.5	0.5 PL	<0.5	<0.5	<0.5	0.5 PL	<0.5	<0.5	<0.5	0.5 PL	<0.5	<0.5	0.5 PL							0.35 - 0.82	NA		
BENZENE	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	0.7 PL	1.0	0.7 PL	0.7 PL	0.7 PL	5.0	2.2	0.7	5.7	5.0	3.1	88						0.6 - 3.3	0.3 - 1.5			
METHYLMETHACRYLATE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	0.5 PL	<0.5	<0.5	0.5 PL	0.5 PL	0.5 PL	13						NA	NA			
METHYL ISOBUTYL KETONE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	0.7						NA	NA			
n-OCTANE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	2.4						0.18 - 0.76	NA		
TOLUENE	0.7	0.5 PL	0.5 PL	0.8	2.0	1.1	0.7	1.2	0.5 PL	0.5 PL	1.3	0.6	0.5 PL	0.5	4.0	2.7	1.2	4.2	3.7	2.5	84							2.16 - 5.21	0.3 - 1.5		
TETRACHLOROETHENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	1.5	0.4 PL	<0.4						0.1 - 0.9	0.1 - 0.5		
n-NONANE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	0.4 PL	0.4 PL	<0.4	3.2						0.14 - 0.81	NA			
CHLOROBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	0.4 PL	0.4 PL	<0.4	1.1						ND - 0.31	<0.4			
ETHYLBENZENE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	0.5 PL	<0.5	0.5 PL	0.5	<0.5	<0.5	<0.5	3.0	0.9	0.6 PL	3.2	2.5	1.8	48							0.2 - 1.2	0.1 - 0.8		
m,p-XYLENE	<0.5	<0.5	<0.5	<0.5	0.5 PL	0.5 PL	0.5 PL	0.5 PL	<0.5	0.5 PL	0.5 PL	0.5 PL	<0.5	<0.5	0.5 PL	<0.5	<0.5	0.5 PL	0.5 PL	0.5 PL	3.2							1.0 - 6.5	0.2 - 1.2		
o-XYLENE	<0.5	<0.5	<0.5	<0.5	<0.5	0.5 PL	<0.5	0.5 PL	<0.5	<0.5	0.5 PL	<0.5	<0.5	<0.5	0.5 PL	<0.5	<0.5	0.5 PL	0.5 PL	0.5 PL	6.5							0.2 - 1.5	0.2 - 1.1		
STYRENE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	0.5 PL	<0.5	<0.5	<0.5	1.9	0.5 PL	<0.5	1.9	1.8	1.1	75							ND - 0.31	<0.24		
ALPHA-PINENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4						0.023 - 0.284	NA			
ISOPROPYLBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	1.1	0.4 PL	<0.4	0.9	0.9	0.5	13							ND	<0.2		
n-DECANE	<0.3	<0.3	<0.3	<0.3	<0.3	0.3 PL	0.3 PL	<0.3	<0.3	<0.3	<0.3	0.3 PL	<0.3	<0.3	0.3 PL	<0.3	<0.3	0.3 PL	0.3 PL	0.3 PL	2.6							0.05 - 0.66	NA		
n-PROPYLBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	<0.4	<0.4	2.5							0.08 - 0.28	<0.2		
1,3,5-TRIMETHYLBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	2.0							0.04 - 0.51	<0.2		
1,2,4-TRIMETHYLBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	0.4 PL	0.4 PL	<0.4	0.4 PL	0.4 PL	<0.4	2.9							0.57 - 1.51	<0.2		
o-LIMONENE	<0.4	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	<0.4	0.4 PL	<0.4	<0.4	<0.4	0.4	0.4	0.4 PL	<0.4	<0.4	<0.4	0.6							NA	NA		
1,2,3-TRIMETHYLBENZENE	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	<0.4	1.4							NA	NA		
1,4-DICHLOROBENZENE	<0.5	<0.5	<0.5	<0.5	<0.5	0.5 PL	1.0	0.5 PL	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5						ND - 0.2	0.1 - 0.6		
n-UNDECANE	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	0.3 PL	<0.3	<0.3	0.3 PL	0.3 PL	<0.3	2.9							ND - 0.62	NA		
n-BUTYLBENZENE	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	1.0							ND	<1.6		
n-DODECANE	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	<0.3	0.3 PL	<0.3	<0.3	0.3 PL	0.3 PL	<0.3	3.7							ND - 0.41	NA		

M - morning (~6-8AM)

A - afternoon (~2-4PM)

E - evening (~6-8PM)

NA - Not Available

ND - Not Detected

Samples were collected by Summa canisters (EPA Method TO-14) for a 2-hour period, the plume sample was a 25 second grab sample. Samples were analyzed by the Wedsworth Center for Labs and Research.

¹The United States Environmental Protection Agency Volatile Organic Compounds Database (EPA) is a compilation of indoor and outdoor data from studies across the United States published in 1988.

²The New York State Department of Health Database (NYSDOH) is a summary of indoor and outdoor air sample results in control homes collected and analyzed by the NYSDOH from 1989 through 1996.

³The levels are the 25th percentile to 75th percentile, (middle half), of the data from the EPA and NYSDOH databases.

< Means less than. The number following this symbol (<) is the lowest level the laboratory test can reliably measure (detection limit). If there is a "<" before any number, then the chemical was NOT detected in the sample.

[PL] - Present, but less than the level indicated.