



Pace Analytical Services, Inc.
1700 Elm Street, Suite 200
Minneapolis, MN 55414
Phone: 612.607.1700
Fax: 612.607.6444

DETERMINATION OF PCDD/PCDF LEVELS

Prepared for:
ATC Associates
Attn: Jim Cohn
104 East 25th Street, 10th Floor
New York, NY 10010



This report contains 23 pages.

The results reported herein conform to the most current NELAC standards, where applicable, unless otherwise narrated in the body of the report.

Project: Chemical Analysis

Client Project Number: 19125-0246

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full, without the written consent of Pace Analytical Services, Inc.



www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES

Pace Analytical Services, Inc.1700 Elm Street, Suite 200
Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE:** December 20, 2001**ISSUED TO:** ATC Associates
Attn: Mr. Jim Cohn
104 East 25th Street, 10th Floor
New York, NY 10010**REPORT NO:** 01-1051682**INTRODUCTION**

This report presents the results from the analyses performed on four samples which were submitted by a representative of ATC Associates. The samples were analyzed for the presence or absence of polychlorinated dibenzo-p-dioxins (PCDDs) and dibenzofurans (PCDFs) using a modified version of USEPA Method 8290 as described below.

SAMPLE IDENTIFICATION

<u>Client ID</u>	<u>Sample Type</u>	<u>Date Received</u>	<u>Pace ID</u>
ATC-1203-STY-DX1	Wipe	12/04/01	103183083
ATC-1203-STY-DX2	Wipe	12/04/01	103183091
ATC-1203-STY-DX3	Wipe	12/04/01	103183117
ATC-1203-STY-DX4	Wipe	12/04/01	103183125

METHODOLOGY**Sample Extraction**

Each sample was spiked with $^{13}\text{C}_{12}$ -labeled PCDD/PCDF internal standards (Table 1) and extracted with toluene in a Soxhlet extractor. The extract was quantitatively transferred to a Kuderna-Danish concentrator, concentrated, and solvent exchanged to hexane. The hexane extract was then spiked with 2,3,7,8-TCDD- $^{37}\text{Cl}_4$ enrichment efficiency standard (Table 1) and processed through the analyte enrichment procedures described below.

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.

**Pace Analytical™**

www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES**Pace Analytical Services, Inc.**

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE: December 20, 2001****PAGE: 2****REPORT NO: 01-1051682****PCDD/PCDF Analyte Enrichment**

The extraction procedure often removes a variety of compounds, in addition to the PCDDs and PCDFs, from the sample matrix. Some of these compounds can directly interfere with the analyses while others can overload the capillary column causing degradation in chromatographic resolution or sensitivity. The analyte enrichment steps described below are used to remove interferences from the extracts.

Each extract was diluted to 100 mL with hexane, transferred to a separatory funnel, and washed with 1N sodium hydroxide, concentrated sulfuric acid, and aqueous sodium chloride (5% w/v) as needed. The hexane extract was quantitatively transferred to a liquid chromatography column containing alternating layers of silica gel, 40% concentrated sulfuric acid on silica gel, and 33% 1 N sodium hydroxide on silica gel. The column was eluted with 90 mL of hexane and the entire eluate was collected and concentrated, under ambient conditions, to a volume of 1 mL.

Each extract was then fractionated on a liquid chromatography column containing 4 g of activated alumina. The column was eluted with 20 mL of hexane followed by 15 mL of 60% methylene chloride/hexane. The 60% methylene chloride/hexane fraction was concentrated to 1 mL under a stream of dry nitrogen and applied to the top of a chromatography column containing 1 g of 5% AX-21 activated carbon in silica gel. The column was eluted with two 2-mL portions of hexane, 2 mL of cyclohexane/methylene chloride (50:50 v/v) and cyclohexane/methanol/toluene (75:20:5 v/v) in the forward direction, and then with toluene in the reverse direction. The toluene fraction was collected, concentrated, spiked with recovery standards (1,2,3,4-TCDD-¹³C₁₂ and 1,2,3,7,8,9-HxCDD-¹³C₁₂) and taken to a final volume of 20 μ L.

PCDD/PCDF Analyses

Each sample extract was analyzed for the presence of PCDDs and PCDFs using combined capillary column gas chromatography/high resolution mass spectrometry (HRGC/HRMS). The instrumentation consisted of a Hewlett Packard Model 6890 gas chromatograph interfaced to a Micromass Ultima high-resolution mass spectrometer. The capillary column was interfaced directly into the ion source of the mass spectrometer, thus providing the highest possible sensitivity while minimizing degradation of the chromatographic resolution.

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.

**Pace Analytical™**

www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES**Pace Analytical Services, Inc.**

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE: December 20, 2001****PAGE: 3****REPORT NO: 01-1051682****PCDD/PCDF Analyses (Cont.)**

The mass spectrometer was operated in the electron impact ionization mode at a mass resolution of 10,000-11,000 ($M/\Delta M$, 10 percent valley definition). This resolution is sufficient to resolve most interferences, such as PCBs, thus providing the highest level of confidence that the detected levels of PCDD/PCDF were not false positives resulting from interferences. Typical operating parameters for the HRGC/HRMS analyses are summarized in Table 2.

The data were acquired by selected-ion-recording (SIR) using groups of ion masses similar to those described in USEPA Method 8290. The five groups corresponded to the tetrachlorinated through octachlorinated congener classes. Each group contained two ion masses for the PCDDs, two ion masses for the PCDFs, the corresponding ion masses from the two isotopically labeled internal standards, and the ion mass characteristic of the polychlorinated diphenylether (PCDE) which, if present, could cause false responses in the dibenzofuran channels.

Each group of ion masses also contained a lock mass which was used by the data system to automatically correct the mass focus of the instrument. The data system determined the centroid of the lock mass during each data acquisition cycle and corrected the mass focus of the analyte and internal standard ion masses to assure that the centers of the mass peaks were being monitored.

The criteria used to judge positive responses for a PCDD/PCDF isomer included:

- * Simultaneous response at both ion masses of the PCDD or PCDF
- * Signal-to-noise ratio equal to or greater than 2.5:1.0 for both ion masses
- * Chlorine isotope ratio within 15% of the theoretical value
- * Chromatographic retention time within +/- 2 seconds of the expected retention time
- * Chromatographic retention times within elution windows determined from analyses of standard mixtures
- * Absence of simultaneous response in the PCDF and PCDE ion traces

A list of the exact ion masses monitored for the determination of PCDD/PCDF isomers and the PCDE interferences is presented in Table 3. Also included are the theoretical chlorine isotope ratios for the ten congener classes.

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.



www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES**Pace Analytical Services, Inc.**1700 Elm Street, Suite 200
Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE: December 20, 2001****PAGE: 4****REPORT NO: 01-1051682****PCDD/PCDF Quantification and Calculations**

The PCDD/PCDF isomers were quantified by comparison of their responses to the responses of the labeled internal standards. Relative response factors were calculated from analyses of standard mixtures containing representatives of each of the PCDD/PCDF congener classes at five concentration levels, and each of the internal standards at one concentration level, as shown in Table 4. The PCDD/PCDF response factors were calculated by comparing the sum of the responses from the two ion masses monitored for each chlorine congener class to the sum of the responses from the two ion masses of the corresponding isotopically labeled internal standard. The formula for the response factor calculation is:

$$R_f = \frac{A_n \times Q_{is}}{A_{is} \times Q_n}$$

where:

R_f = Response factor
 A_n = Sum of integrated areas for native isomer
 Q_{is} = Quantity of labeled internal standard
 A_{is} = Sum of integrated areas for labeled internal standard
 Q_n = Quantity of native isomer

The levels of PCDD/PCDF in each sample were quantified using the following equation:


$$C = \frac{A_n \times Q_{is}}{A_{is} \times W \times R_f}$$

where:

C = Concentration of target isomer or congener class
 A_n = Sum of integrated areas for the target isomer or congener class
 Q_{is} = Quantity of labeled internal standard added to the sample
 A_{is} = Sum of integrated areas for the labeled internal standard
 W = Sample amount
 R_f = Response factor

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES**Pace Analytical Services, Inc.**

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE: December 20, 2001****PAGE: 5****REPORT NO: 01-1051682****PCDD/PCDF Quantification and Calculations (Cont.)**

Each pair of ion mass peaks in the selected-ion-current chromatograms was evaluated manually to determine if it met the criteria for a PCDD or PCDF isomer. Areas of all peaks exhibiting correct ion ratios, having retention times within the correct windows, and having areas corresponding to concentrations in the range covered by the initial calibration were then summed for calculations of total congener concentrations. The toxic equivalence of each sample was calculated using the factors listed in Table 5.

A limit of detection (LOD) based on producing a signal that is 2.5 times the noise level, was calculated for each undetected 2,3,7,8-substituted isomer of any tetra through octa chlorinated congener class. The noise heights used to calculate the detection limits were measured at the retention time of the specific isomer. The formula used for calculating the LOD is:

$$\text{LOD} = \frac{H_n \times Q_{is} \times 2.5}{H_{is} \times W \times R_f}$$

where:

LOD = Single isomer limit of detection
 H_n = Sum of noise heights at native isomer retention time
 Q_{is} = Quantity of labeled internal standard
 H_{is} = Sum of peak heights for labeled internal standard
 W = Sample amount
 R_f = Response factor

The recovery of the 2,3,7,8-TCDD- $^{37}\text{Cl}_4$ enrichment efficiency standard and each $^{13}\text{C}_{12}$ -labeled internal standard, relative to either 1,2,3,4-TCDD- $^{13}\text{C}_{12}$ or 1,2,3,7,8,9-HxCDD- $^{13}\text{C}_{12}$, was calculated using the following equation:

$$\%R = \frac{A_{is} \times Q_{rs} \times 100\%}{R_{fr} \times A_{rs} \times Q_{is}}$$

where:

$\%R$ = Percent recovery of labeled internal standard
 A_{is} = Sum of integrated areas of labeled internal standard
 Q_{rs} = Quantity of recovery standard
 A_{rs} = Sum of integrated areas of recovery standard
 R_{fr} = Response factor of the specific labeled internal standard relative to the recovery standard
 Q_{is} = Quantity of the labeled internal standard congener added to the sample

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
 without the written consent of Pace Analytical Services, Inc.



Pace Analytical Services, Inc.
1700 Elm Street, Suite 200
Minneapolis, MN 55414
Phone: 612.607.1700
Fax: 612.607.6444

REPORT OF: CHEMICAL ANALYSES

PROJECT: PCDD/PCDF ANALYSES

DATE: December 20, 2001

PAGE: 6

REPORT NO: 01-1051682

Quality Control for PCDD/PCDF Analyses

The performance of the sample processing steps and the instrumentation are monitored on a routine basis. The procedures and criteria are summarized below.

One method blank and one laboratory spike sample are typically prepared with each ten samples of any given matrix. Recoveries of the native PCDD/PCDF analytes in the laboratory spike samples generally range from 70 to 130%. Recoveries of selected analytes outside this range do not invalidate the data but provide information, which is used by the laboratory to monitor recovery trends and to assure optimization of the method.

Internal standards are spiked into each sample prior to extraction in order to monitor the level of recovery, which is achieved for each individual sample. Acceptable recoveries range from 40 to 135 percent for the internal standards unless a deviation is due to variation in instrument response as a result of analytical interferences.

The resolution of the mass spectrometer is verified prior to each analysis to be 10,000 or greater. Hardcopies of the reference peaks are printed at the beginning and end of each analysis day. The resolving power of the DB-5MS chromatographic column is checked daily by analyzing a standard solution containing 2,3,7,8-TCDD and the adjacent TCDD isomers. The DB-225 column resolution is checked daily by analyzing a standard solution containing 2,3,7,8-TCDF and the adjacent TCDF isomers. Acceptable performance is achieved when 2,3,7,8-TCDD or 2,3,7,8-TCDF is resolved from the adjacent isomers by a valley of 25% or less. The group times for the selected-ion-monitoring data acquisitions are also checked daily by analyzing the column performance mix which has been modified to contain the first and last eluting isomers of each congener class. In this way one is assured of collecting data representative of the total PCDD/PCDF content and that the 2,3,7,8-substituted isomers are suitably resolved.

Initial calibrations are generated by analyzing standard solutions (see Table 4) containing target native and labeled PCDD/PCDF compounds. Response factors are calculated and averaged for each compound. These averages are used for quantification and for comparison to the daily continuing calibration. The relative standard deviation for each native compound must be 20% or less (30% or less for the labeled compounds) as specified in Method 8290. A continuing calibration standard is analyzed at the beginning and end of each 12-hour shift on days when initial calibrations are not performed. The initial calibration is considered to be valid when the response factors from the continuing calibration analysis fall to within the ranges specified in Method 8290.

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.

**Pace Analytical™**

www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES**Pace Analytical Services, Inc.**

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES**DATE: December 20, 2001****PAGE: 7****REPORT NO: 01-1051682****RESULTS**

The results from the analyses are presented in the following:

Appendix A - Documentation

Appendix B - PCDD/PCDF Analysis Results

DISCUSSION


The recoveries of the isotopically-labeled PCDD/PCDF internal standards in the sample extracts ranged from 26-117% and indicate a level of efficiency through the extraction and enrichment steps that is considered typical to somewhat low for this matrix. With the exception of two standards, all of the internal standard recoveries for the field samples were within the target limits for this method. Also, since the quantifications of the native 2,3,7,8-substituted isomers were based on isotope dilution, the data were automatically corrected for variation in recovery and accurate values were obtained.

A laboratory method blank was prepared and analyzed with the sample batch as part of our routine quality control procedures. The results, found at the beginning of Appendix B, show the blank to contain a trace level of non-2,3,7,8 substituted TCDD. Two of the samples contained this isomer at a level similar to that seen in the blank and are flagged "BJ" on the data summary sheets. In general, levels less than ten times the background are not considered statistically different from the background.

A laboratory spike sample was also prepared with the sample batch by extracting a clean wipe that had been fortified with native standard materials. The results, found at the end of Appendix B, show that the spiked native compounds were recovered at 80-110%. This indicates a high degree of accuracy for these determinations.

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.



Pace Analytical™

www.pacelabs.com

REPORT OF: CHEMICAL ANALYSES

Pace Analytical Services, Inc.

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

PROJECT: PCDD/PCDF ANALYSES

DATE: December 20, 2001


PAGE: 8

REPORT NO: 01-1051682


REMARKS

The sample extracts will be retained for a period of 30 days from the date of this report and then discarded unless other arrangements are made. The raw mass spectral data will be archived on magnetic tape for a period of not less than one year. Questions regarding the data contained in this report may be directed to the authors at the numbers provided below.

Pace Analytical Services, Inc.




Charles V. Sueper, Technical Director
High Resolution Mass Spectrometry
(612) 607-6387



Scott C. Unze
Project Manager, Dioxins
(612) 607-6383

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

www.pacelabs.com

Pace Analytical Services, Inc.

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700


Fax: 612.607.6444

TABLE 1. Spike Levels of PCDD/PCDF Standards

Internal Standards	Spike Level (ng)
2,3,7,8-TCDF- ¹³ C ₁₂	2.0
2,3,7,8-TCDD- ¹³ C ₁₂	2.0
1,2,3,7,8-PeCDF- ¹³ C ₁₂	2.0
2,3,4,7,8-PeCDF- ¹³ C ₁₂	2.0
1,2,3,7,8-PeCDD- ¹³ C ₁₂	2.0
1,2,3,4,7,8-HxCDF- ¹³ C ₁₂	2.0
1,2,3,6,7,8-HxCDF- ¹³ C ₁₂	2.0
1,2,3,7,8,9-HxCDF- ¹³ C ₁₂	2.0
2,3,4,6,7,8-HxCDF- ¹³ C ₁₂	2.0
1,2,3,4,7,8-HxCDD- ¹³ C ₁₂	2.0
1,2,3,6,7,8-HxCDD- ¹³ C ₁₂	2.0
1,2,3,4,6,7,8-HpCDF- ¹³ C ₁₂	2.0
1,2,3,4,7,8,9-HpCDF- ¹³ C ₁₂	2.0
1,2,3,4,6,7,8-HpCDD- ¹³ C ₁₂	2.0
OCDD- ¹³ C ₁₂	4.0
<u>Recovery Standards</u>	
1,2,3,4-TCDD- ¹³ C ₁₂	2.0
1,2,3,7,8,9-HxCDD- ¹³ C ₁₂	2.0
<u>Enrichment Efficiency Standard</u>	
2,3,7,8-TCDD- ³⁷ Cl ₄	0.2

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.

**Pace Analytical™**

www.pacelabs.com

Pace Analytical Services, Inc.

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700


Fax: 612.607.6444

**TABLE 2. High Resolution PCDD/PCDF Analyses
HRGC/HRMS Operating Parameters**

Mass Resolution	10,000-11,000 (M/ Δ M, 10% valley)
Electron Energy	32 electron volts
Accelerating Voltage	8,000 volts
Source Temperature	275°C
Preamplifier Gain	10 ⁻⁶ amp/volt
Multiplier Gain	~10 ⁵
Chromatographic Column	60 M DB-5MS
Transfer Line Temperature	260°C
Injection Mode	Splitless
Carrier Gas	Helium
Carrier Flow Velocity	~30 cm/sec

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

www.pacelabs.com

Pace Analytical Services, Inc.

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

**TABLE 3. Exact Ion Masses Monitored
for the Determination of PCDDs, PCDFs, and PCDEs**

Ratio Compound	Accurate Mass		Theoretical
	Mass 1	Mass 2	Mass 1/Mass 2
Tetra-CDDs	319.8965	321.8936	0.77
Tetra-CDFs	303.9016	305.8987	0.77
Hexa-CDPEs	375.8364		
Penta-CDDs	355.8546	357.8517	1.54
Penta-CDFs	339.8597	341.8567	1.54
Hepta-CDPEs	409.7974		
Hexa-CDDs	389.8156	391.8127	1.23
Hexa-CDFs	373.8207	375.8178	1.23
Octa-CDPEs	445.7555		
Hepta-CDDs	423.7766	425.7737	1.03
Hepta-CDFs	407.7817	409.7788	1.03
Nona-CDPEs	479.7165		
Octa-CDD	457.7377	459.7347	0.88
Octa-CDF	441.7428	443.7398	0.88
Deca-CDPE	513.6775		


CDDs = Chlorinated Dibenzo-p-dioxins

CDFs = Chlorinated Dibenzofurans

CDPEs = Chlorinated Diphenylethers

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

www.pacelabs.com

Pace Analytical Services, Inc.1700 Elm Street, Suite 200
Minneapolis, MN 55414

Phone: 612.607.1700


Fax: 612.607.6444

TABLE 4. High Resolution Calibration Solutions

Native CDDs/CDFs	CS1	Concentration (pg/uL)			
		CS2	CS3	CS4	CS5
2,3,7,8-TCDD	0.5	2	10	40	200
2,3,7,8 TCDF	0.5	2	10	40	200
1,2,3,7,8-PeCDD	2.5	10	50	200	1000
1,2,3,7,8-PeCDF	2.5	10	50	200	1000
2,3,4,7,8-PeCDF	2.5	10	50	200	1000
1,2,3,4,7,8-HxCDD	2.5	10	50	200	1000
1,2,3,6,7,8-HxCDD	2.5	10	50	200	1000
1,2,3,7,8,9-HxCDD	2.5	10	50	200	1000
1,2,3,4,7,8-HxCDF	2.5	10	50	200	1000
1,2,3,6,7,8-HxCDF	2.5	10	50	200	1000
1,2,3,7,8,9-HxCDF	2.5	10	50	200	1000
2,3,4,6,7,8-HxCDF	2.5	10	50	200	1000
1,2,3,4,6,7,8-HpCDD	2.5	10	50	200	1000
1,2,3,4,6,7,8-HpCDF	2.5	10	50	200	1000
1,2,3,4,7,8,9-HpCDF	2.5	10	50	200	1000
OCDD	5.0	20	100	400	2000
OCDF	5.0	20	100	400	2000
Internal Standards					
2,3,7,8-TCDD- ¹³ C ₁₂	100	100	100	100	100
2,3,7,8-TCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,7,8-PeCDD- ¹³ C ₁₂	100	100	100	100	100
1,2,3,7,8-PeCDF- ¹³ C ₁₂	100	100	100	100	100
2,3,4,7,8-PeCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,4,7,8-HxCDD- ¹³ C ₁₂	100	100	100	100	100
1,2,3,6,7,8-HxCDD- ¹³ C ₁₂	100	100	100	100	100
1,2,3,4,7,8-HxCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,6,7,8-HxCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,7,8,9-HxCDF- ¹³ C ₁₂	100	100	100	100	100
2,3,4,6,7,8-HxCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,4,6,7,8-HpCDD- ¹³ C ₁₂	100	100	100	100	100
1,2,3,4,6,7,8-HpCDF- ¹³ C ₁₂	100	100	100	100	100
1,2,3,4,7,8,9-HpCDF- ¹³ C ₁₂	100	100	100	100	100
OCDD- ¹³ C ₁₂	200	200	200	200	200
Recovery Standards					
1,2,3,4-TCDD- ¹³ C ₁₂	100	100	100	100	100
1,2,3,7,8,9-HxCDD- ¹³ C ₁₂	100	100	100	100	100
Enrichment Efficiency Standard					
2,3,7,8-TCDD- ³⁷ C ₁₄	0.5	2	10	40	200

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

www.pacelabs.com

Pace Analytical Services, Inc.

1700 Elm Street, Suite 200

Minneapolis, MN 55414

Phone: 612.607.1700

Fax: 612.607.6444

**TABLE 5. 2,3,7,8-TCDD Equivalency Factors (TEFs) for the
Polychlorinated Dibenzo-p-dioxins and Dibenzofurans**

Number	Compound(s)	TEF
1	2,3,7,8-TCDD	1.00
2	1,2,3,7,8-PeCDD	0.50
3	1,2,3,6,7,8-HxCDD	0.1
4	1,2,3,7,8,9-HxCDD	0.1
5	1,2,3,4,7,8-HxCDD	0.1
6	1,2,3,4,6,7,8-HpCDD	0.01
7	OCDD	0.001
8	* Total - TCDD	0.0
9	* Total - PeCDD	0.0
10	* Total - HxCDD	0.0
11	* Total - HpCDD	0.0
12	2,3,7,8-TCDF	0.10
13	1,2,3,7,8-PeCDF	0.05
14	2,3,4,7,8-PeCDF	0.5
15	1,2,3,6,7,8-HxCDF	0.1
16	1,2,3,7,8,9-HxCDF	0.1
17	1,2,3,4,7,8-HxCDF	0.1
18	2,3,4,6,7,8-HxCDF	0.1
19	1,2,3,4,6,7,8-HpCDF	0.01
20	1,2,3,4,7,8,9-HpCDF	0.01
21	OCDF	0.001
22	* Total - TCDF	0.0
23	* Total - PeCDF	0.0
24	* Total - HxCDF	0.0
25	* Total - HpCDF	0.0

*Excluding the 2,3,7,8-substituted congeners.

Reference: 1989 ITEFs

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.



Pace Analytical Services, Inc.
1700 Elm Street, Suite 200
Minneapolis, MN 55414
Phone: 612.607.1700
Fax: 612.607.6444

Appendix A

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full;
without the written consent of Pace Analytical Services, Inc.

Pace Analytical
www.paceanalytical.com

CHAIN-OF-CUSTODY / Analytical Request Document

The Chain-of-Custody is a LEGAL DOCUMENT. All relevant fields must be completed accurately.

Page: 1 of 1

Required Client Information: Section A

Company: **ATC Associates**
Address: **104 E 25th Street**
New York NY 10010
Phone: **212/353-2222** Fax: **212/353-2306**

Required Client Information: Section B

Report To: **James Cohn**
Copy To:
Invoice To:
P.O.
Project Name: **Hyvessant HS**
Project Number: **1925-0246**

Client Information (Check quote/contract):

Requested Due Date:
TAT:
TAT in calendar days less than 14 days subject
Inquiry and contractual obligations end may
In a 24 Hour Turnaround Surcharge.
Turnaround Time (TAT) in calendar days.

To Be Completed by Pace Analytical Client: Section C

Quote Reference:
Project Manager:
Project #: **1051682**
Profile #:

Required Client Information: Section D

SAMPLE ID (One character per box.)

Valid Matrix Codes
Matrix Code
SWT
BL
CL
WV
AP
TJ
OT

ITEM NUMBER	Required Client Information: Section D										Vial Matrix Codes		DATE COLLECTED	VOLUME COLLECTED	Preservatives								Requested Analysis										REMARKS / Lab ID
	SAMPLE ID (One character per box.)										Matrix Code				Unpreserved	H2SO4	HNO3	HCl	NaOH	Na2S2O8	Methanol	Dioxin											
1	A	C	I	O	-	T	-	X			WP	12/03	1 m2																			03183083	
2	A	C	I	O	-	T	-	X			WP	12/03	1 m2																			091	
3	A	C	I	O	-	T	-	X			WP	12/03	1 m2																			117	
4	A	C	I	O	-	T	-	X			WP	12/03	Blank																			125	
5																																	
6																																	
7																																	
8																																	
9																																	
10																																	
11																																	
12																																	

SHIPMENT METHOD	ARRIVAL NO.	SHIPMENT DATE	NO. OF SAMPLES	TIME	REQUESTED BY / AFFILIATION	DATE	TIME	ACCEPTED BY / AFFILIATION	DATE	TIME
-----------------	-------------	---------------	----------------	------	----------------------------	------	------	---------------------------	------	------

SAMPLE CONDITION: SAMPLE NOTES:

Temp in C: **10°C**
Received on Ice: **Y/N**
Stored Cooler: **Y/N**
Sample Intact: **Y/N**

all samples area
1 m2
(box)

SAMPLE NAME AND ADDRESS

PRINT Name of SAMPLER:

WAGDZ

SIGNATURE of SAMPLER

[Signature]

DATE: Signed: **12/03/01**



Pace Analytical Services, Inc.
1700 Elm Street, Suite 200
Minneapolis, MN 55414
Phone: 612.607.1700
Fax: 612.607.6444

Appendix B

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace AnalyticalTM

Pace Analytical Services, Inc.
1700 Elm Street - Suite 200
Minneapolis, MN 55414

Tel: 612-607-1700
Fax: 612-607-6444

Method 8290 Blank Analysis Results

Client - ATC

Lab Sample ID	BLANK-1358	Matrix	WIPE
Filename	F11208A_04	Dilution	NA
Total Amount Extracted	1 Sample	Extracted	12/05/2001
ICAL Date	11/19/2001	Analyzed	12/08/2001 19:46
CCal Filename(s)	F11208A_01 & F11208A_17	Injected By	BAL

Native Isomers	Conc ng/S	EMPC ng/S	LRL ng/S	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	ND	----	0.00200	2,3,7,8-TCDF-13C	2.00	62
Total TCDF	ND	----	0.00200	2,3,7,8-TCDD-13C	2.00	73
				1,2,3,7,8-PeCDF-13C	2.00	75
2,3,7,8-TCDD	ND	----	0.00200	2,3,4,7,8-PeCDF-13C	2.00	75
Total TCDD	0.0033	----	0.00200 J	1,2,3,7,8-PeCDD-13C	2.00	85
				1,2,3,4,7,8-HxCDF-13C	2.00	87
1,2,3,7,8-PeCDF	ND	----	0.01000	1,2,3,6,7,8-HxCDF-13C	2.00	88
2,3,4,7,8-PeCDF	ND	----	0.01000	2,3,4,6,7,8-HxCDF-13C	2.00	86
Total PeCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDF-13C	2.00	77
				1,2,3,4,7,8-HxCDD-13C	2.00	77
1,2,3,7,8-PeCDD	ND	----	0.01000	1,2,3,6,7,8-HxCDD-13C	2.00	95
Total PeCDD	ND	----	0.01000	1,2,3,4,6,7,8-HpCDF-13C	2.00	80
				1,2,3,4,7,8,9-HpCDF-13C	2.00	63
1,2,3,4,7,8-HxCDF	ND	----	0.01000	1,2,3,4,6,7,8-HpCDD-13C	2.00	73
1,2,3,6,7,8-HxCDF	ND	----	0.01000	OCDD-13C	4.00	47
2,3,4,6,7,8-HxCDF	ND	----	0.01000			
1,2,3,7,8,9-HxCDF	ND	----	0.01000	1,2,3,4-TCDD-13C	2.00	NA
Total HxCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	ND	----	0.01000	2,3,7,8-TCDD-37Cl4	0.20	76
1,2,3,6,7,8-HxCDD	ND	----	0.01000			
1,2,3,7,8,9-HxCDD	ND	----	0.01000			
Total HxCDD	ND	----	0.01000			
1,2,3,4,6,7,8-HpCDF	ND	----	0.01000	Total 2,3,7,8-TCDD		
1,2,3,4,7,8,9-HpCDF	ND	----	0.01000	Equivalence: 0.00 ng/S		
Total HpCDF	ND	----	0.01000	(Using ITE Factors)		
1,2,3,4,6,7,8-HpCDD	ND	----	0.01000			
Total HpCDD	ND	----	0.01000			
OCDF	ND	----	0.02000			
OCDD	ND	----	0.02000			

Conc = Concentration (Totals include 2,3,7,8-substituted isomers).
EMPC = Estimated Maximum Possible Concentration
LRL = Lower Reporting Limit
J = Concentration detected is below the calibration range
P = Recovery outside of target range
A = Detection Limit based on signal-to-noise measurement

I = Interference
E = PCDE Interference
ND = Not Detected
NA = Not Applicable
NC = Not Calculated
* = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.



Pace Analytical Services, Inc.
1700 Elm Street - Suite 200
Minneapolis, MN 55414

Tel: 612-607-1700
Fax: 612-607-6444

Method 8290 Analysis Results

Client - ATC

Client's Sample ID	ATC-1203-STY-DX1		
Lab Sample ID	103183083		
Filename	U11209A_12		
Injected By	BAL		
Total Amount Extracted	1 Sample	Matrix	WIPE
% Moisture	NA	Dilution	NA
Dry Weight Extracted	NA	Collected	12/03/2001
ICAL Date	11/10/2001	Received	12/04/2001
CCal Filename(s)	U11209A_01 & U11209A_17	Extracted	12/05/2001
Method Blank ID	BLANK-1358	Analyzed	12/10/2001 04:30

Native Isomers	Conc ng/S	EMPC ng/S	LRL ng/S	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	ND	----	0.00200	2,3,7,8-TCDF-13C	2.00	82
Total TCDF	0.0022	----	0.00200 J	2,3,7,8-TCDD-13C	2.00	76
				1,2,3,7,8-PeCDF-13C	2.00	79
2,3,7,8-TCDD	ND	----	0.00200	2,3,4,7,8-PeCDF-13C	2.00	79
Total TCDD	0.0044	----	0.00200 BJ	1,2,3,7,8-PeCDD-13C	2.00	78
				1,2,3,4,7,8-HxCDF-13C	2.00	112
1,2,3,7,8-PeCDF	ND	----	0.01000	1,2,3,6,7,8-HxCDF-13C	2.00	94
2,3,4,7,8-PeCDF	ND	----	0.01000	2,3,4,6,7,8-HxCDF-13C	2.00	104
Total PeCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDF-13C	2.00	97
				1,2,3,4,7,8-HxCDD-13C	2.00	94
1,2,3,7,8-PeCDD	ND	----	0.01000	1,2,3,6,7,8-HxCDD-13C	2.00	108
Total PeCDD	0.0150	----	0.01000 J	1,2,3,4,6,7,8-HpCDF-13C	2.00	86
				1,2,3,4,7,8,9-HpCDF-13C	2.00	75
1,2,3,4,7,8-HxCDF	ND	----	0.01000	1,2,3,4,6,7,8-HpCDD-13C	2.00	74
1,2,3,6,7,8-HxCDF	ND	----	0.01000	OCDD-13C	4.00	48
2,3,4,6,7,8-HxCDF	ND	----	0.01000			
1,2,3,7,8,9-HxCDF	ND	----	0.01000	1,2,3,4-TCDD-13C	2.00	NA
Total HxCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	ND	----	0.01000	2,3,7,8-TCDD-37Cl4	0.20	74
1,2,3,6,7,8-HxCDD	ND	----	0.01000			
1,2,3,7,8,9-HxCDD	ND	----	0.01000			
Total HxCDD	0.0380	----	0.01000 J			
1,2,3,4,6,7,8-HpCDF	ND	----	0.01000	Total 2,3,7,8-TCDD		
1,2,3,4,7,8,9-HpCDF	ND	----	0.01000	Equivalence: 0.00016 ng/S		
Total HpCDF	ND	----	0.01000	(Using ITE Factors)		
1,2,3,4,6,7,8-HpCDD	0.0100	----	0.01000 J			
Total HpCDD	0.0100	----	0.01000 J			
OCDF	ND	----	0.02000			
OCDD	0.0560	----	0.02000 J			


Conc = Concentration (Totals include 2,3,7,8-substituted isomers)
EMPC = Estimated Maximum Possible Concentration
A = Detection Limit based on signal-to-noise measurement
J = Concentration detected is below the calibration range
B = Less than 10 times higher than method blank level
P = Recovery outside of target range
Nn = Value obtained from additional analysis

LRL = Lower Reporting Limit
I = Interference
E = PCDE Interference
S = Saturated signal
ND = Not Detected
NA = Not Applicable
NC = Not Calculated
* = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

Pace Analytical Services, Inc.
1700 Elm Street - Suite 200
Minneapolis, MN 55414

Tel: 612-607-1700
Fax: 612-607-6444

Method 8290 Analysis Results

Client - ATC

Client's Sample ID	ATC-1203-STY-DX2		
Lab Sample ID	103183091		
Filename	U11209A_13		
Injected By	BAL		
Total Amount Extracted	1 Sample	Matrix	WIPE
% Moisture	NA	Dilution	NA
Dry Weight Extracted	NA	Collected	12/03/2001
ICAL Date	11/10/2001	Received	12/04/2001
CCal Filename(s)	U11209A_01 & U11209A_17	Extracted	12/05/2001
Method Blank ID	BLANK-1358	Analyzed	12/10/2001 05:21

Native Isomers	Conc ng/S	EMPC ng/S	LRL ng/S	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	ND	----	0.00200	2,3,7,8-TCDF-13C	2.00	80
Total TCDF	ND	----	0.00200	2,3,7,8-TCDD-13C	2.00	76
				1,2,3,7,8-PeCDF-13C	2.00	85
2,3,7,8-TCDD	ND	----	0.00200	2,3,4,7,8-PeCDF-13C	2.00	84
Total TCDD	0.0089	----	0.00200 BJ	1,2,3,7,8-PeCDD-13C	2.00	86
				1,2,3,4,7,8-HxCDF-13C	2.00	117
1,2,3,7,8-PeCDF	ND	----	0.01000	1,2,3,6,7,8-HxCDF-13C	2.00	106
2,3,4,7,8-PeCDF	ND	----	0.01000	2,3,4,6,7,8-HxCDF-13C	2.00	108
Total PeCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDF-13C	2.00	95
				1,2,3,4,7,8-HxCDD-13C	2.00	101
1,2,3,7,8-PeCDD	ND	----	0.01000	1,2,3,6,7,8-HxCDD-13C	2.00	109
Total PeCDD	ND	----	0.01000	1,2,3,4,6,7,8-HpCDF-13C	2.00	88
				1,2,3,4,7,8,9-HpCDF-13C	2.00	71
1,2,3,4,7,8-HxCDF	ND	----	0.01000	1,2,3,4,6,7,8-HpCDD-13C	2.00	73
1,2,3,6,7,8-HxCDF	ND	----	0.01000	OCDD-13C	4.00	48
2,3,4,6,7,8-HxCDF	ND	----	0.01000			
1,2,3,7,8,9-HxCDF	ND	----	0.01000	1,2,3,4-TCDD-13C	2.00	NA
Total HxCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	ND	----	0.01000	2,3,7,8-TCDD-37Cl4	0.20	73
1,2,3,6,7,8-HxCDD	ND	----	0.01000			
1,2,3,7,8,9-HxCDD	ND	----	0.01000			
Total HxCDD	0.0120	----	0.01000 J			
1,2,3,4,6,7,8-HpCDF	ND	----	0.01000	Total 2,3,7,8-TCDD		
1,2,3,4,7,8,9-HpCDF	ND	----	0.01000	Equivalence: 0.000065 ng/S		
Total HpCDF	ND	----	0.01000	(Using ITE Factors)		
1,2,3,4,6,7,8-HpCDD	ND	----	0.01000			
Total HpCDD	ND	----	0.01000			
OCDF	ND	----	0.02000			
OCDD	0.0650	----	0.02000 J			

Conc = Concentration (Totals include 2,3,7,8-substituted isomers)
EMPC = Estimated Maximum Possible Concentration
A = Detection Limit based on signal-to-noise measurement
J = Concentration detected is below the calibration range
B = Less than 10 times higher than method blank level
P = Recovery outside of target range
Nn = Value obtained from additional analysis

LRL = Lower Reporting Limit
I = Interference
E = PCDE Interference
S = Saturated signal
ND = Not Detected
NA = Not Applicable
NC = Not Calculated
* = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace AnalyticalTM

 Pace Analytical Services, Inc.
 1700 Elm Street - Suite 200
 Minneapolis, MN 55414

 Tel: 612-607-1700
 Fax: 612- 607-6444

Method 8290 Analysis Results

Client - ATC

Client's Sample ID	ATC-1203-STY-DX3	Matrix	WIPE
Lab Sample ID	103183117	Dilution	NA
Filename	U11209A_14	Collected	12/03/2001
Injected By	BAL	Received	12/04/2001
Total Amount Extracted	1 Sample	Extracted	12/05/2001
% Moisture	NA	Analyzed	12/10/2001 06:12
Dry Weight Extracted	NA		
ICAL Date	11/10/2001		
CCal Filename(s)	U11209A_01 & U11209A_17		
Method Blank ID	BLANK-1358		

Native Isomers	Conc ng/S	EMPC ng/S	LRL ng/S	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	0.0041	----	0.00200 J	2,3,7,8-TCDF-13C	2.00	80
Total TCDF	0.0670	----	0.00200	2,3,7,8-TCDD-13C	2.00	75
				1,2,3,7,8-PeCDF-13C	2.00	78
2,3,7,8-TCDD	ND	----	0.00200	2,3,4,7,8-PeCDF-13C	2.00	76
Total TCDD	0.1100	----	0.00200	1,2,3,7,8-PeCDD-13C	2.00	79
				1,2,3,4,7,8-HxCDF-13C	2.00	117
1,2,3,7,8-PeCDF	ND	----	0.01000	1,2,3,6,7,8-HxCDF-13C	2.00	101
2,3,4,7,8-PeCDF	ND	----	0.01000	2,3,4,6,7,8-HxCDF-13C	2.00	106
Total PeCDF	0.0130	----	0.01000 J	1,2,3,7,8,9-HxCDF-13C	2.00	94
				1,2,3,4,7,8-HxCDD-13C	2.00	99
1,2,3,7,8-PeCDD	ND	----	0.01000	1,2,3,6,7,8-HxCDD-13C	2.00	106
Total PeCDD	0.1200	----	0.01000	1,2,3,4,6,7,8-HpCDF-13C	2.00	82
				1,2,3,4,7,8,9-HpCDF-13C	2.00	63
1,2,3,4,7,8-HxCDF	ND	----	0.01000	1,2,3,4,6,7,8-HpCDD-13C	2.00	66
1,2,3,6,7,8-HxCDF	ND	----	0.01000	OCDD-13C	4.00	39 P
2,3,4,6,7,8-HxCDF	ND	----	0.01000			
1,2,3,7,8,9-HxCDF	ND	----	0.01000	1,2,3,4-TCDD-13C	2.00	NA
Total HxCDF	0.0310	----	0.01000 J	1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	ND	----	0.01000	2,3,7,8-TCDD-37Cl4	0.20	75
1,2,3,6,7,8-HxCDD	ND	----	0.01000			
1,2,3,7,8,9-HxCDD	ND	----	0.01000			
Total HxCDD	0.1200	----	0.01000			
1,2,3,4,6,7,8-HpCDF	0.0290	----	0.01000 J	Total 2,3,7,8-TCDD		
1,2,3,4,7,8,9-HpCDF	ND	----	0.01000	Equivalence: 0.0016 ng/S		
Total HpCDF	0.0290	----	0.01000 J	(Using ITE Factors)		
1,2,3,4,6,7,8-HpCDD	0.0520	----	0.01000			
Total HpCDD	0.0790	----	0.01000			
OCDF	0.0260	----	0.02000 J			
OCDD	0.3400	----	0.02000			

Conc = Concentration (Totals include 2,3,7,8-substituted isomers)

EMPC = Estimated Maximum Possible Concentration

A = Detection Limit based on signal-to-noise measurement

J = Concentration detected is below the calibration range

B = Less than 10 times higher than method blank level

P = Recovery outside of target range

Nn = Value obtained from additional analysis

LRL = Lower Reporting Limit

I = Interference

E = PCDE Interference

S = Saturated signal

ND = Not Detected

NA = Not Applicable


NC = Not Calculated

* = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

 This report shall not be reproduced, except in full,
 without the written consent of Pace Analytical Services, Inc.


Pace Analytical™

Pace Analytical Services, Inc.
1700 Elm Street - Suite 200
Minneapolis, MN 55414

Tel: 612-607-1700
Fax: 612-607-6444

Method 8290 Analysis Results

Client - ATC

Client's Sample ID	ATC-1203-STY-DX4		
Lab Sample ID	103183125		
Filename	U11209A_15		
Injected By	BAL		
Total Amount Extracted	1 Sample	Matrix	WIPE
% Moisture	NA	Dilution	NA
Dry Weight Extracted	NA	Collected	12/03/2001
ICAL Date	11/10/2001	Received	12/04/2001
CCal Filename(s)	U11209A_01 & U11209A_17	Extracted	12/05/2001
Method Blank ID	BLANK-1358	Analyzed	12/10/2001 07:03

Native Isomers	Conc ng/S	EMPC ng/S	LRL ng/S	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	ND	----	0.00200	2,3,7,8-TCDF-13C	2.00	86
Total TCDF	ND	----	0.00200	2,3,7,8-TCDD-13C	2.00	78
				1,2,3,7,8-PeCDF-13C	2.00	84
2,3,7,8-TCDD	ND	----	0.00200	2,3,4,7,8-PeCDF-13C	2.00	83
Total TCDD	ND	----	0.00200	1,2,3,7,8-PeCDD-13C	2.00	84
				1,2,3,4,7,8-HxCDF-13C	2.00	109
1,2,3,7,8-PeCDF	ND	----	0.01000	1,2,3,6,7,8-HxCDF-13C	2.00	101
2,3,4,7,8-PeCDF	ND	----	0.01000	2,3,4,6,7,8-HxCDF-13C	2.00	103
Total PeCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDF-13C	2.00	93
				1,2,3,4,7,8-HxCDD-13C	2.00	98
1,2,3,7,8-PeCDD	ND	----	0.01000	1,2,3,6,7,8-HxCDD-13C	2.00	107
Total PeCDD	ND	----	0.01000	1,2,3,4,6,7,8-HpCDF-13C	2.00	82
				1,2,3,4,7,8,9-HpCDF-13C	2.00	61
1,2,3,4,7,8-HxCDF	ND	----	0.01000	1,2,3,4,6,7,8-HpCDD-13C	2.00	68
1,2,3,6,7,8-HxCDF	ND	----	0.01000	OCDD-13C	4.00	26 IP
2,3,4,6,7,8-HxCDF	ND	----	0.01000			
1,2,3,7,8,9-HxCDF	ND	----	0.01000	1,2,3,4-TCDD-13C	2.00	NA
Total HxCDF	ND	----	0.01000	1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	ND	----	0.01000	2,3,7,8-TCDD-37Cl4	0.20	73
1,2,3,6,7,8-HxCDD	ND	----	0.01000			
1,2,3,7,8,9-HxCDD	ND	----	0.01000			
Total HxCDD	ND	----	0.01000			
1,2,3,4,6,7,8-HpCDF	ND	----	0.01000	Total 2,3,7,8-TCDD		
1,2,3,4,7,8,9-HpCDF	ND	----	0.01000	Equivalence: 0.000036 ng/S		
Total HpCDF	ND	----	0.01000	(Using ITE Factors)		
1,2,3,4,6,7,8-HpCDD	ND	----	0.01000			
Total HpCDD	ND	----	0.01000			
OCDF	ND	----	0.02000			
OCDD	0.036	----	0.02000 J			


Conc = Concentration (Totals include 2,3,7,8-substituted isomers)
 EMPC = Estimated Maximum Possible Concentration
 A = Detection Limit based on signal-to-noise measurement
 J = Concentration detected is below the calibration range
 B = Less than 10 times higher than method blank level
 P = Recovery outside of target range
 Nn = Value obtained from additional analysis

LRL = Lower Reporting Limit
 I = Interference
 E = PCDE Interference
 S = Saturated signal
 ND = Not Detected
 NA = Not Applicable
 NC = Not Calculated
 * = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.


Pace AnalyticalTM

Pace Analytical Services, Inc.
1700 Elm Street - Suite 200
Minneapolis, MN 55414

Tel: 612-607-1700
Fax: 612-607-6444

Method 8290 Laboratory Control Spike Results

Client - ATC

Lab Sample ID	SPIKE-1362	Matrix	WIPE
Filename	U11209A_02	Dilution	NA
Total Amount Extracted	1 Sample	Extracted	12/05/2001
ICAL Date	11/10/2001	Analyzed	12/09/2001 20:04
CCal Filename(s)	U11209A_01 & U11209A_17	Injected By	BAL
Method Blank ID	BLANK-1358		

Native Isomers	Qs (ng)	Qm (ng)	% Rec.	Internal Standards	ng's Added	Percent Recovery
2,3,7,8-TCDF	0.20	0.16	81	2,3,7,8-TCDF-13C	2.00	73
				2,3,7,8-TCDD-13C	2.00	73
				1,2,3,7,8-PeCDF-13C	2.00	67
2,3,7,8-TCDD	0.20	0.16	80	2,3,4,7,8-PeCDF-13C	2.00	65
				1,2,3,7,8-PeCDD-13C	2.00	71
				1,2,3,4,7,8-HxCDF-13C	2.00	90
1,2,3,7,8-PeCDF	1.00	0.96	96	1,2,3,6,7,8-HxCDF-13C	2.00	80
2,3,4,7,8-PeCDF	1.00	0.97	97	2,3,4,6,7,8-HxCDF-13C	2.00	81
				1,2,3,7,8,9-HxCDF-13C	2.00	81
				1,2,3,4,7,8-HxCDD-13C	2.00	101
1,2,3,7,8-PeCDD	1.00	0.96	96	1,2,3,6,7,8-HxCDD-13C	2.00	85
				1,2,3,4,6,7,8-HpCDF-13C	2.00	80
				1,2,3,4,7,8,9-HpCDF-13C	2.00	75
1,2,3,4,7,8-HxCDF	1.00	0.96	96	1,2,3,4,6,7,8-HpCDD-13C	2.00	85
1,2,3,6,7,8-HxCDF	1.00	0.96	96	OCDD-13C	4.00	61
2,3,4,6,7,8-HxCDF	1.00	0.96	96			
1,2,3,7,8,9-HxCDF	1.00	0.95	95	1,2,3,4-TCDD-13C	2.00	NA
				1,2,3,7,8,9-HxCDD-13C	2.00	NA
1,2,3,4,7,8-HxCDD	1.00	1.00	100	2,3,7,8-TCDD-37Cl4	0.20	71
1,2,3,6,7,8-HxCDD	1.00	0.96	96			
1,2,3,7,8,9-HxCDD	1.00	0.93	93			
1,2,3,4,6,7,8-HpCDF	1.00	1.00	100			
1,2,3,4,7,8,9-HpCDF	1.00	0.99	99			
1,2,3,4,6,7,8-HpCDD	1.00	0.96	96			
OCDF	2.00	1.74	87			
OCDD	2.00	2.20	110			

Qs = Quantity Spiked
Qm = Quantity Measured
Rec. = Recovery (Expressed as Percent)
P = Recovery outside of target range
X = Background subtracted value
Nn = Value obtained from additional analysis
NA = Not Applicable
* = See Discussion

Report No.....01-1051682

REPORT OF LABORATORY ANALYSIS

This report shall not be reproduced, except in full,
without the written consent of Pace Analytical Services, Inc.