NYC 9/11 Public Portal Document

ATC ASSOCIATES INC.

104 East 25th St., 10th Floo
New York, New York 1001(
www.atc-enviro.com
212.353.828(
Fax 212.353.830(

January 16, 2002

Mr. Bernard P. Orlan
Director, Environmental Health and Safety
New York City Board of Education
28-11 Queens Plaza North
Long Island City, NY 11101

Re: Air Sampling for Metals
Stuyvesant High School
ATC Project No. 15-19125-0246

Dear Mr. Orlan:

ATC Associates, Inc. (ATC) is pleased to provide this letter report of air sampling for metals in airborne dust on January 9, 2002, at the referenced school. The New York City Board of Education (BOE) requested this sampling at the recommendation of H. A. Bader, a consultant retained by the parents association.

Scope of Work

ATC collected three (3) samples for metals at specified locations. Samples were collected for eight (8) hours using 37 mm mixed cellulose ester filter connected to battery-powered sampling pumps at a flow rate of 2 liters per minute. Pumps were calibrated before and after each sampling period. Two (2) samples were collected inside the building and one (1) was collected on the roof. One blank sample was also submitted on each day. The samples were sent to PSC Analytical Services, Inc., an American Industrial Hygiene Association-accredited laboratory in Reading, PA. Analysis was performed by NIOSH Methods 7300M. Each sample was analyzed for a panel of eleven (11) metals. The metals were aluminum (Al), cadmium (Cd), chromium (Cr), cobalt (Co), copper (Cu), iron (Fe), magnesium (Mg), manganese (Mn), molybdenum (Mo), nickel (Ni) and zinc (Zn). A copy of the analytical report is attached.

The sampling locations, specified by the parents association consultant, were as follows:

- 1st floor lobby;
- 2nd floor bridge entrance hallway;
- Roof by the air intake.

The established regulatory values for metals are intended for industrial workplace exposures and are not generally applicable to office/school settings. There are no standards for these metals in indoor air in non-industrial situations. The Environmental Protection Agency has set a screening level for hexavalent Cr (the more toxic form) of 0.6 micrograms per cubic meter of air (0.6 ug/m^3). These samples were analyzed for both hexavalent and trivalent Cr. The EPA level is based on continuous exposure for one year.

CONFIDENTIAL, CITY_21MC100_003429292

SUBJECT TO PROTECTIVE ORDER NO. 2, SEPT. 2008

NYC-WTC_000238167

NYC 9/11 Public Portal Document

Results

The levels of airborne metals did not exceed the limits of detection in any of the samples.

ATC appreciates the opportunity to provide this indoor air quality service to the BOE. Please call me at (212) 353-8280 extension 212 if you have any questions.

Sincerely,
ATC Associates, Inc.

James R. Cohn, CIH, CSP
Regional Manager
Industrial Hygiene Division

Attachment

NYC-WTC_000238168